arXiv:0711.2881v1 [astro-ph] 19 Nov 2007

FULL TITLE
*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***
NAMES OF EDITORS

The Milky Way is an Exceptionally Quiet galaxy: Implications for spiral formation

M. Puech[1,2], F. Hammer[2], L. Chemin[2], H. Flores[2], and M. Lehnert[2]

[1] *ESO, Karl-Schwarzschild-Strasse 2, D-85748 Garching bei München, Germany*
[2] *GEPI, Observatoire de Paris, CNRS, University Paris Diderot; 5 Place Jules Janssen, 92190 Meudon, France*

Abstract. We compare both the Milky Way and M31 to local external disk galaxies within the same mass range, using their relative locations in the planes formed by V_{flat} vs. M_K (the Tully-Fisher relation), j_{disk} (specific angular momentum) and the average Fe abundance of stars in the galaxy outskirts. We find, for all relationships, that the MW is systematically offset by 1 σ or more, showing a significant deficiency in stellar mass, angular momentum, disk radius and [Fe/H] in the stars in its outskirts at a given V_{flat}. Our Galaxy appears to have escaped any significant merger over the last 10-11 Gyr which may explain its peculiar properties. As with M31, most local spirals show evidence for a history shaped mainly by relatively recent merging.

1. The MW is a very peculiar galaxy

In Fig. 1, we show the local Tully-Fisher Relation (TFR) in K-band, as well as the specific angular momentum of the disk j_{disk} vs. the rotational velocity V_{flat}. In both relations, V_{flat} is taken as a proxy for the total mass. Noteworthy, the local TFR was carefully derived comparing three different local samples (see top-left inset and Hammer et al. 2007). Strikingly, the MW systematically falls at $\sim 1\,\sigma$ of these relations, conversely to M31: the MW has a smaller stellar mass and specific angular momentum, compared to local galaxies of similar total mass. On the basis of their location in the (M_K, V_{flat}, R_d) volume, the fraction of spirals like the MW is ∼7%, while M31 appears to be a much more typical spiral (see Hammer et al. 2007). The representativity of the MW as a typical spiral galaxy is actually even smaller, if one takes into account the iron abundance of the stellar halo (see right panel).

2. Implication for spiral formation

One can explain the differences between the MW and M31 by the absence of significant merger events over the last 10-11 Gyr for the former. On the contrary, M31 has been much more influenced by mergers (and possibly a major merger) during the last 8 Gyrs (see Hammer et al. 2007 for a detailed discussion). Could the bulk of local spirals be shaped and assembled through a rich past merger history during the last 8 Gyrs? Such a scenario is indeed suggested by direct observations of violent dynamical processes at work in z∼0.6 galaxies



Figure 1. *Left and middle panels:* K-band TFR and j_{disk} vs. V_{flat} relation for three local galaxy samples (blue triangles from Verheijen 2001, red dots from Pizagno et al. 2007, and black squares from Courteau 1997). Solid lines show the fits for the combined three samples, and the dashed lines the residual 1 σ dispersion. The locations of the MW and M31 are marked by two large green dots. In the top-left panel insert, we show the individual fits of the TF relations for each of the three samples (see more details in Hammer et al. 2007). In the middle panel, open stars show results of simulations by Steinmetz & Navarro (1999), illustrating the difficulty in reproducing the angular momentum of spiral galaxies when using the standard model for disk formation. *Right panel:* Iron abundances of stellar haloes of eight spiral galaxies (from Mouhcine et al. 2006) vs. log(V_{flat}) (small red points). Large green points mark the position of the Milky Way and M31. The full line assumes $M_{star} \sim V^4$ (McGaugh 2005) and $M_{star} \sim Z^2$ following the prescription of Dekel & Silk (1987). Red dashed lines identify a box including the color measurements of Zibetti et al. (2004), after stacking 1047 edge-on SDSS galaxies, and assuming that their colors are dominated by red giant stars.

(Yang et al. 2007), and by the relatively important major merger rate per M* galaxy during the last 11 Gyrs (0.5-0.7 up to z$\sim$1, see Hammer et al. 2007, and 4-5 up to z$\sim$3, see Conselice et al. 2003). In this context, rich major merger episodes in the past may explain the Tully-Fisher relation, the angular momentum problem and the abundances of stars in spiral outskirts. What remains unclear is whether or not these properties could be explained by the standard scenario for spiral formation, i.e., a monolithic collapse followed by pure secular evolution.

Acknowledgments. We wish to thank the organizing committees for this very pleasant meeting.

References

Conselice, C.J., Gallagher, J.S. III, & Wyse, R.F.G. 2003, AJ, 125, 66
Courteau, S. 1997, AJ, 114, 2402
Dekel, A., & Silk, J. 1986, ApJ, 303, 39
Hammer, F., Flores, H., Elbaz, D., Zheng, X. Z., Liang, Y. C., & Cesarsky, C. 2005, A&A, 430, 115
Hammer, F., Puech, M., Chemin, L., Flores, H., & Lehnert, M. 2007, ApJ, 662, 322
McGaugh, S.S. 2005, ApJ, 632, 859
Mouhcine, M., Bamford, S.P., Aragón-Salamanca, A., et al. 2006, MNRAS, 369, 891
Pizagno, J., Prada, F., Weinberg, D.H., et al. 2007, AJ, 134, 945
Steinmetz, M. & Navarro, J. 1999, ApJ, 513, 555
Verheijen, M.A.W. 2001, ApJ, 563, 694
Yang, Y., Flores, H., Hammer, F., et al. 2007, A&A, accepted

Zibetti, S., White, S.D.M., & Brinchmann, J. 2004, MNRAS, 347, 556